

**AFRL-AFOSR-UK-TR-2010-0010**



**Correlated Measurements of the Disturbances of  
Geomagnetic Field and Changes of Secondary Particle Fluxes  
at Aragats-Space Environmental Center (ASEC)**

**Ashot A. Chilingarian**

**Alikhanyan Physics Institute  
Cosmic Ray Division  
2 Alikhanian Brothers St.  
Yerevan, Armenia 37036**

**EOARD GRANT 09-3053**

**February 2010**

**Final Report for 1 January 2009 to 31 December 2009**

**Distribution Statement A: Approved for public release distribution is unlimited.**

**Air Force Research Laboratory  
Air Force Office of Scientific Research  
European Office of Aerospace Research and Development  
Unit 4515 Box 14, APO AE 09421**

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p><b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b></p>					
1. REPORT DATE (DD-MM-YYYY) 24-02-2010		2. REPORT TYPE Final Report		3. DATES COVERED (From – To) 1 January 2009 – 31 December 2009	
4. TITLE AND SUBTITLE  Correlated Measurements of the Disturbances of Geomagnetic Field and Changes of Secondary Particle Fluxes at Aragats-Space Environmental Center (ASEC)			5a. CONTRACT NUMBER FA8655-09-1-3053		
			5b. GRANT NUMBER  Grant 09-3053		
			5c. PROGRAM ELEMENT NUMBER  61102F		
			5d. PROJECT NUMBER		
6. AUTHOR(S)  Dr. Ashot A Chilingarian			5d. TASK NUMBER		
			5e. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Alikhanyan Physics Institute 2 Alikhanian Brothers St. Yerevan 37036 Armenia			8. PERFORMING ORGANIZATION REPORT NUMBER  Grant 09-3053		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  EOARD Unit 4515 BOX 14 APO AE 09421			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)  AFRL-AFOSR-UK-TR-2010-0010		
12. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
<p><b>14. ABSTRACT</b> Our objective is the development of a geomagnetic storms forecast service. It will consist of a observed solar bursts data, particle detector data and current geomagnetic activity data. They originate in solar activity and variability of many different forms including solar flares, coronal mass ejections, solar wind sector boundaries and coronal hole streams. Variations of geomagnetic field can adversely affect any technical systems which rely on local magnetic field measurements to be taken in the absence of significant geomagnetic variations. Users who are unaware of and thus unprepared for the imminent occurrence of major magnetic field disturbances may conduct operations which later turn out to have been useless and eventually a waste of time and resources. Our project attempts to address geomagnetic storm forecasting by developing a service which gives advance notice of increasing geomagnetic activity and thus enables the affected users to plan and conduct their operations in a more cost-effective way. Information on flux changes for different secondary particles helps to test Interplanetary Magnetic Fields (IMF) and magnetosphere for understanding of the level of disturbance and specific mechanisms leading to cutoff rigidity reduction. Our observations of the GMS occurred during 23rd Solar activity cycle demonstrate that Cosmic Ray increase during GMS occurs coherently (or ~ 1 hour in advance) with abrupt changes of the geomagnetic field, sized up by Dst index. To test geomagnetic disturbances models and provide a valuable user-oriented forecast, it is not sufficient to have only global geomagnetic indices. Several studies demonstrate that local geomagnetic indices are better correlated with consequences of severe GMS. Unfortunately in the radius of 1000 km. from mt. Aragats there are no permanently operating geomagnetic observatories or magneto-telluric stations. In the framework of the project we equip ASEC with modern facilities allowing precise measuring geomagnetic and electric fields at location of ASEC monitors. Special low-noise magnetometric laboratory was setup at premises of the Nor Amberd research station of Yerevan Physics Institute at altitude of 2000 m. The LEMI-417 magnetometer was purchased from Lviv Space Research Center of Ukrainian Space agency. The device was installed and tuned with help of Ukrainian experts. After testing and comparison with world data and analogical device purchased later the data of geomagnetic field first time in Armenia is monitored and stored in ASEC data bases.</p> <p>Conclusion - The new facilities for precise measuring of the geomagnetic field are installed and giving data, available on-line. Data will be used for the forecasting of the upcoming geomagnetic field along with data on Solar bursts and secondary cosmic ray data. The magnetometer operates stable, noise level is low.</p>					
15. SUBJECT TERMS EOARD, Space Science					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES  12	19a. NAME OF RESPONSIBLE PERSON SCOTT DUDLEY, Lt Col, USAF
a. REPORT UNCLAS	b. ABSTRACT UNCLAS	c. THIS PAGE UNCLAS			19b. TELEPHONE NUMBER (Include area code) +44 (0)1895 616162

**Project FA8655-09-1-3053**

***Correlated measurements of the disturbances of geomagnetic field and changes of secondary particle fluxes at Aragats-Space Environmental Center (ASEC)***

**Final Report January 1, 2009 – December 31 2009**

**Summary**

Our objective is the development of a geomagnetic storms forecast service. It will consist of a observed solar bursts data, particle detector data and current geomagnetic activity data.

They originate in solar activity and variability of many different forms including solar flares, coronal mass ejections, solar wind sector boundaries and coronal hole streams. Variations of geomagnetic field can adversely affect any technical systems which rely on local magnetic field measurements to be taken in the absence of significant geomagnetic variations. Users who are unaware of and thus unprepared for the imminent occurrence of major magnetic field disturbances may conduct operations which later turn out to have been useless and eventually a waste of time and resources.

Our project attempts to address geomagnetic storm forecasting by developing a service which gives advance notice of increasing geomagnetic activity and thus enables the affected users to plan and conduct their operations in a more cost-effective way.

Information on flux changes for different secondary particles helps to “test” Interplanetary Magnetic Fields (IMF) and magnetosphere for understanding of the level of disturbance and specific mechanisms leading to cutoff rigidity reduction.

Our observations of the GMS occurred during 23<sup>rd</sup> Solar activity cycle demonstrate that Cosmic Ray increase during GMS occurs coherently (or ~ 1 hour in advance) with abrupt changes of the geomagnetic field, sized up by Dst index.

To test geomagnetic disturbances models and provide a valuable user-oriented forecast, it is not sufficient to have only global geomagnetic indices. Several studies demonstrate that local geomagnetic indices are better correlated with consequences of severe GMS. Unfortunately in the radius of 1000 km. from mt. Aragats there are no permanently operating geomagnetic observatories or magneto-telluric stations. In the framework of the project we equip ASEC with modern facilities allowing precise measuring geomagnetic and electric fields at location of ASEC monitors.

Special low-noise magnetometric laboratory was setup at premises of the Nor Amberd research station of Yerevan Physics Institute at altitude of 2000 m. The LEMI-417 magnetometer was purchased from Lviv Space Research Center of Ukrainian Space agency. The device was installed and tuned with help of Ukrainian experts. After testing and comparison with world data and analogical device purchased later the data of geomagnetic field first time in Armenia is monitored and stored in ASEC data bases.

## Introduction

The Earth's magnetic field varies both in space and time. Historically, magnetic observatories were established to monitor the secular change of the Earth's magnetic field, and this remains one of their most important functions. Over 70 countries operate more than 200 observatories worldwide. The magnetic observatory mean data are crucial to the studies of secular change, investigations into the Earth's interior, and to global modeling efforts.

Huge magnetized plasma clouds and shocks initiated by Coronal Mass Ejections (CME) traveling in the interplanetary space with velocities up to 2000 Km/sec (so called Interplanetary Coronal Mass Ejection (ICME)), are known as major drivers of severe space weather conditions. On its way to Earth ICME disturb the IMF. These disturbances “modulate” flux of Galactic Cosmic Ray (GCR) introducing anisotropy and changing energy (rigidity) spectra of the previously isotropic population of protons and stripped nuclei accelerated in the numerous galactic sources. Changes of rather stable flux of GCR are detected by space-born spectrometers (rigidities up to  $\sim 1\text{GV}$ ) and by world-wide networks of particle detectors (rigidities up to  $\sim 100\text{GV}$ ) located at different latitudes, longitudes and altitudes. Therefore, changes of GCR fluxes (so called precursors) can be used for forewarning on upcoming ICME and, consequently on expected GMS.

Most geoeffective types of ICMEs, so called magnetic clouds (cylindrical flux rope structures with helical magnetic field lines, see for details (Mulligan and Russell, 2001) are directly connected to the flaring region of Sun where Solar Cosmic Rays (SCR) are accelerated. Recently “flux-rope” like structures were observed by the payload of the THEMIS satellite constellation ([http://www.nasa.gov/mission\\_pages/themis/news/](http://www.nasa.gov/mission_pages/themis/news/)). This can explain the “collisionless” transport of SCR via “highways” inside the magnetic system connecting Sun with ICME. The cascade particles generated by these SCR with GV rigidities can be registered by surface detectors during so called Ground Level Enhancement (GLE). Fast arrival of the solar particles of highest energies can alert on upcoming abundant SCR flux (of much more higher MeV energies) dangerous for satellite electronics and some surface industries.

ICME is a major modulating agent, forming depletion and enhancement regions of GCR, manifested themselves as peaks and deeps in time series of surface particle detectors. Both GCR and SCR can be used for “probing” ICME, providing highly cost-effective information on the key characteristics of the interplanetary disturbances. Because cosmic rays are fast and have large scattering mean free paths in the solar wind, this information travels rapidly and may provide useful for space weather forecasting (Leerungrat et al., 2003).

Size and strength of the “frozen” magnetic field in an ICME are correlated with the modulation effects ICME poses during its propagation up to 1 AU on the energy spectra and direction of the GCR. In the same time the presence of strong and long duration southward magnetic field is the primary requirement of ICME geoeffectiveness (Valtonen, 2007). Thus, strong magnet field “frozen” in ICME is both modulation agent of GCR and driver of GMS.

The magnetic field found in the magnetic cloud of ICME usually has a well-formed flux-rope structure (see Koskinen and Huttunen, 2006, also see cross-section of the magnetic “rope” - a twisted bundle of magnetic fields) connecting Earth's upper atmosphere directly to the Sun, observed by the THEMIS satellites on May 20, 2007 (NASA science, 20.03.08). At November 20, 2003 the magnetic cloud was directly connected with solar eruptive region and southward magnetic field was enhanced by an order of magnitude (up to  $\sim 50\text{ nT}$ ).

It is not possible to assert that there is one-to-one correspondence between variations of the GCR and strength of GMS, furthermore, besides IMCEs, there are other drivers of storms and modulation agents of GCR. However, large  $B_z$  value associated with approaching ICME is best known diagnostics of GMS strength. Therefore, as was mentioned in the statistical study (Kudela & Storini, 2006) appropriate observations of the variations of the primary and secondary cosmic rays can be a proxy of  $B_z$  value available long before IMCE reach libration point at 1.5 mln. km from Earth where  $B_z$  can be measured directly.

Direct detection of the Energetic Storm Particles (ESP) by the Electron, Proton, and Alpha Monitor (EPAM) instrument on board the Advanced Composition Explorer (ACE) space station (Gold, 1998) can alerts hours prior to the approaching interplanetary shock and plasma cloud (ACE news, 2003). Another space station located in libration point 1.5 mln kilometers from Earth - Solar and Heliospheric Observatory (SOHO) - detected relativistic electrons by its Comprehensive Suprathermal and Energetic Particle Analyzer (COSTEP) instrument. Enhancements in the electron flux also can point on approaching ICME. The modulation effects posed by ICME on the particles of higher energies, not measurable by space-born facilities due to very weak fluxes, are detected by the world-wide networks of Neutron Monitors (NM, responding to GCR energies  $\geq 10$  GeV) and Muon Telescopes (MT, responding to GCR energies  $\geq 15$  GeV) well before the onset of a major geomagnetic storm (Belov et al., 2003, Munakata et al., 2000).

To conclude, ICMEs dominate intense storm occurrences and simultaneously they are correlated with variations of spectra of particles and nuclei in the interplanetary space, ranging from isothermal solar wind ions till GeV energy galactic cosmic rays. Therefore, for developing timely and reliable forecasting service it is necessary to get more insight in the correlations of IMCE parameters with geospace parameters, including changing intensities of the particle fluxes measured at the Earth surface.

## **Instrumentation**

To study the Earth's magnetic field the best instrument is the magnetotelluric station LEMI-417 which was specially developed for the long-term monitoring of geomagnetic field. Construction of device ensure thermal and temporal stability and low noise level for magnetic measurements. The high-sensitive magnetic sensor of flux-gate type, which mainly determines these parameters, was manufactured using well-proved technology on the base of marble and quartz combination implementing recent findings in the excitation circuit construction. The waterproof sensor has the support with bubble level to provide its proper leveling at the selected place (see Fig.1).



**Figure 1 Magnetotelluric station LEMI-417 set (left) and its front panel (right)**

Its main technical parameters are given in the Table 2 below:

Measured range of total magnetic field	$\pm 65000$ nT
Resolution along each component both at the display and registered into the internal FLASH-memory	0.01 nT
Temperature drift	$<0.2$ nT/°C
Frequency band	DC-0.3 Hz
Magnetometer output noise in frequency band (0.03 – 1)Hz	$< 10$ pT rms
Magnetic sensor components orthogonality error	$<30$ min of arc
Automated offset compensation band along each magnetic component	$\pm 65000$ nT
Noise of electric meter in the frequency band 0.03 - 1 Hz	$<0.5$ $\mu$ V rms
Sample rate	4 per s
Volume of the internal FLASH-memory	2 GB
Digital output	RS-232
GPS timing and coordinates determination	
Operating temperature range	Minus 10 to +50°C
Temperature sensors (both in magnetic sensor and electronic units) resolution	0.1 °C
Power supply	$+12^{+6}_{-3}$ V
Power consumption	$<0.9$ W
Weight:	1.1 kg

## Device installation and testing

Magnetometric station LEMI-417, commissioned by Lviv center of Space Research Institute of Ukrainian Academy of Science, was installed in Nor Amberd, 2000 m above sea level (the same type of station with added electrical field sensors will be installed at Aragats, 3200 above sea level). One-minute time series of the 3-dimensional measurements of the geomagnetic field enter the data base of the Aragats Space Environmental Center (ASEC) and will highly improve research of correlations of the geomagnetic parameters and changes of the fluxes of cosmic rays now underway at Aragats. ASEC measures neutral and charged fluxes of secondary cosmic rays (Chilingarian et.al., 2005); correlations of changes of these fluxes with geomagnetic parameters and IMF parameters measured by facilities of the ACE space station located 1.5 mln. km from Earth will assist to understand Solar influence on the Earth environments (see, for instance Bostanjyan & Chilingarian, 2009, Chilingarian & Bostanjyan, 2009). Measurements of geomagnetic field from Nor Amberd will support forewarning of the upcoming major geomagnetic storm. Alert service planned to be provided by CRD in late 2010.




**Figure 2 CRD PhD student Tigran Karapetyan examining geomagnetic field disturbances**



**Figure 3 CRD Postdoc Artur Reymers establishing Internet connection of the geomagnetic laboratory to CRD headquarters. Each minute measurement of the 3 components of geomagnetic field will be send to CRD servers for multivariate analysis and alert services.**


Operation of magnetometric station started in July 2009. In Figure 4 we post the pattern of the first Geomagnetic Storm detected in Nor Amberd in compoarison with DST index (geomagnetic field disturbances averaged other several magnetometers located in middle-low latitudes). Geomagnetic activity increased to major storm levels from 0300 - 1200 UTC on 22 July. Activity decreased to quiet levels at all latitudes on 26 July. ACE solar wind measurements indicated the storm conditions on 22 July were associated with the onset of a coronal hole high-speed wind stream. Solar wind velocities began to gradually increase early on 22 July and eventually reached a maximum of 601 km/sec at 24/0158 UTC. Interplanetary magnetic field (IMF) changes associated with the onset of the high-speed stream included an increase in IMF B<sub>t</sub> (peak 18 nT at 22/0533 UTC) and a sustained period of southward IMF B<sub>z</sub> (minimum -18 nT at 22/0703 UTC).





**Figure 4 Comparison of the geomagnetic storm detected by new facility in Nor Amberd and DST index calculated by Kyoto magnetic observatory (DST is measured in relative units; Nor Amberd magnetometer – in absolute)**

The device provide possibility to measure the intensity of the magnetic field each second, usually we integrate the intensity above 1 minute interval to be compatible with particle detectors. However, the short variations of the geomagnetic field also can be studied. After purchasing another magnetometer of the same type we provide comparison studies of both detectors. Second magnetometer was located in the main building of the Nor Amberd station and was affected by the noise posed by movement of personnel and cars. It is why the pattern is not identical (see Figure 5) and shifted. However the overall closeness of both time series is apparent.



**Figure 5 Comparison of the  $B_z$  measurements taken by 2 same type magnetometers located at magnetometric lab and Nor Amherd main building**

## Conclusion

The new facilities for precise measuring of the geomagnetic field are installed and giving data, available on-line. Data will be used for the forecasting of the upcoming geomagnetic field along with data on Solar bursts and secondary cosmic ray data. The magnetometer operates stable, noise level is low.

## Reference

- Chilingarian, A., et al., Correlated measurements of secondary cosmic ray fluxes by the Aragats Space – Environmental Center monitors, Nucl. Instrum. Methods Phys. Res., A543, 483-496, 2005.
- Chilingarian, A., and N. Bostanjyan, Cosmic ray intensity increases detected by Aragats Space Environmental Center monitors during the 23rd solar activity cycle in correlation with geomagnetic storms, J. Geophys. Res., doi:10.1029/2009JA014346, 2009.
- Chilingarian, A., Bostanjyan, N., On the Relation of the Forbush Decreases Detected by ASEC Monitors during the 23rd Solar Activity Cycle with ICME Parameters, Advances in Space Research, doi: 10.1016/j.asr.2009.09.00, 2009
- Mulligan T., and Russell C.T. (2001), Multispacecraft modeling of the flux rope structure of interplanetary coronal mass ejections, J. Geophys. Res., 106, 10581-10596.
- K. Leerunnavarat , D. Ruffolo, J. W. Bieber (2003), The Astrophysical Journal, 593, 587–596.
- Koskinen H.E.J., Huttunen K.M.J., Space Weather: From Solar Eruptions to Magnetospheric Storms, in Solar Eruptions and Energetic Particles, Geophysical Monographs Series 165, 375-385, 2006
- Kudela, K. and Storini, M.: Possible tools for space weather issues from cosmic ray continuous records, Advances in Space Research, Volume 37, Issue 8, 1443-1449, 2006.
- R. E. Gold, S. M. Krimigis, et al., (1998), Space Sci. Rev., 86, 541.
- ACE News , 2004, ACE News #86 – Nov 14, 2004 Real-time Solar Wind Data from ACE make Accurate Forecasts of Magnetic Storms a Reality,
- Belov, A.V., Bieber, J.W., Eroshenko, E.A., et al, (2003) Adv. Space Res., 31, No 4, 919-924.
- Munakata, K., Bieber, J.W., Yasue, S., et al.: Precursors of geomagnetic storms observed by the muon detector network, J. Geophys. Res. 105, 27457-27468, 2000.
- N. Gopalswamy, S. Yashiro, and S. Akiyama «Geoeffectiveness of halo coronal mass ejections» JOURNAL OF GEOPHYSICAL RESEARCH, VOL. 112, A06112, doi:10.1029/2006JA012149, 2007,

## ACRONIMS

CE	Advanced Composition Explorer
ADAS	Advanced Data Analysis System
ADC	Amplitude-Digital-Converters
AMMM	Aragats Multidirectional Muon Monitor
ANM	Aragats Neutron Monitor
ASCII	American Standard Code for Information Interchange
ASEC	Aragats Space Environmental Center
ASNT	Aragats Solar Neutron Telescope
AU	Astronomical unit (distance Sun-Earth, $1,5 \times 10^{11}$ m)
CME	Coronal Mass Ejection
CMOS	Complementary Metal–Oxide–Semiconductor
CPLD	Complex Programmable Logic Device
CR	Cosmic Rays
CRD	Cosmic Ray Division, Yerevan Physics Institute
DAQ	Data Acquisition
DVIN	Data Visualization Interactive Network EAS Extensive Air Showers
ESA	European Space Agency
Fd	Forbush decrease
GCR	Galactic Cosmic Rays
GLE	Ground level Enhancement (Excess, Event)
GMS	Geomagnetic Storm;
GOES	Geostationary Operational Environmental Satellite
GPS	Global Positioning System
ICME	Interplanetary Coronal Mass Ejection;
IHY	International Heliophysical Year
IMF	Interplanetary Magnetic Field IP Interplanetary
LADC	Logarithmic ADC
LAN	Local Area Network
NAMMM	Nor Amberd Multidirectional Muon Monitor
NANM	Nor-Amberd Neutron Monitor NM Neutron Monitor
PMT	Photo Multiplier Tube
SCR	Solar Cosmic Ray
SEP	Solar Energetic Particles
SEVAN	Space Environmental Viewing and Analysis Network
SF	Solar Flare
SNT	Solar Neutron Telescopes
SVG	Scalable Vector Graphics
TTL	Transistor-Transistor Logic
URCS	Unified Readout and Control Server
XML	eXtensible Markup Language